

# How does fungal diversity change based on woody debris type? A case study in Northern Spain

Nerea Abrego\*,

Isabel Salcedo

Dept. Plant Biology and Ecology (Botany),  
University of the Basque Country (UPV/EHU),  
P. D. Box 644, E-48080 Bilbao, Spain

Studies on lignicolous fungal communities in southern Europe are scarce and most of them focused on coarse woody debris (CWD). Fine (FWD) and very fine woody debris (VFWD) are rarely studied. For this study, all woody debris of 90 beechwood plots (in Navarre, Northern Spain) was checked to search for lignicolous macromycetes. Although most species were found on CWD, small sized woody debris being especially diverse on Ascomycota species, and big sized woody debris on Basidiomycota ones. The CAP analysis revealed that fungal communities from different diameter woody debris classes were dissimilar and that fungal composition changed gradually, from smaller diameter woody debris towards coarser ones, and vice versa. Formed groups were also different in fungal species richness, VFWD and CWD groups having a higher number of habitual and frequent species. Additionally, fungal composition changed according to beechwood types; thus faithful species for each woody debris type in different beechwood types were found. Finally, the structure and composition of fungal community changed in different decay stages and the CAP analysis showed three main groups: communities associated with weakly, moderately and strongly decayed wood.

**Key words:** decay stage, diameter, *Fagus sylvatica*, lignicolous fungal community, macromycetes, Navarre

## INTRODUCTION

Dead wood is an essential component in forest ecosystems, maintaining carbon store in long term and providing the nutrients and habitat to a large number of organisms, including birds, mammals, arthropods, nematodes, bryophytes, lichens, fungi and bacteria. Fungi are a crucial part of this biodiversity (Siitonen, 2001), and they play a key role for the diversity of organisms associated with dead wood. The enzymatic and physical actions of wood-inhabiting fungi chemically and structurally change the wood resulting in numerous habitats and food resources (Niemelä et al., 1995; Renvall, 1995) that are irreplaceable for most of the other organisms living in dead wood (Boddy, 2001).

Nevertheless, organisms associated with wood represent an important but vulnerable group (Heilmann-Clausen, Christensen, 2004). Man has severely manipulated the European beech forest landscape for centuries (Peterken, 1996) causing a simplification of biological and structural complexity in most present-day beech forests. In this way, the regular extraction of any woody debris in traditional forest management has caused a large decrease of this habitat (Harmon et al., 1986). In addition, forest fragmentation has imposed further difficulties for dispersal of dead wood dependent organisms (Lonsdale et al., 2008). The combination of these reasons has caused most wood dependent organisms to have decreased or become extinct locally and regionally (ECCB 1995; Siitonen, 2001).

\* Corresponding author. E-mail: nerea.abrego@ehu.es

At the regional scale, climatic conditions, soil type, tree species composition, continuity and management history have been identified as important variables influencing fungal species diversity and community structure (Boddy, 1992; Sippola, Renvall, 1999; Küffer, Senn-Irlet, 2005a; Heilmann-Clausen et al., 2005). At the local scale, wood inhabiting fungal composition is conditioned by dead wood quality, which is mostly influenced by three features: size of dead woody debris, degree of decomposition and host tree species (Küffer, 2008). Most of the works that have studied wood inhabiting fungal diversity have focused on large logs and trunks (Bader et al., 1995; Heilmann-Clausen, Christensen, 2003), although recently the emphasis has been laid on smaller woody debris (Heilmann-Clausen, Christensen, 2004; Nordén et al., 2004; Küffer, Senn-Irlet, 2005b). Results have shown that branches and twigs may be particularly rich in species where there is little other substrate available, such as in managed forests (Küffer, Senn-Irlet, 2005b), and that ascomycetes are especially highly dependent on fine woody debris (Nordén et al., 2004).

As dead wood undergoes physical and chemical changes during the decomposition process, a successional pathway is established, firstly colonising pioneer species, later being replaced by others. Some studies have revealed that decay stage is the most important variable for understanding fungal community at the local scale (Renvall, 1995; Høiland, Bendiksen, 1996; Heilmann-Clausen, 2001).

In Northern and Central Europe several studies about the effect of wood inhabiting fungi on the biodiversity of forest ecosystems have been carried out, however, in Southern Europe there are still areas where lignicolous fungal diversity is not fully known (Tellería, 2002) and studies of lignicolous fungal communities are scarce, such as the Iberian Peninsula (Heilmann-Clausen, Walley, 2007). It is also noteworthy that there are few studies that have dealt with all the macromycetes fruiting in dead wood, especially not taking into account those that grow on very fine woody debris and resupinate fungi due to their laborious collection and identification, despite being the main agents of wood decay (Swift, 1982).

This study aims to answer the following questions: (1) How diverse and dissimilar are the fungal communities in different-sized woody debris? (2) Is it possible to assess the changes in the fungal community throughout the wood decaying process?

## MATERIALS AND METHODS

### Study area

The area of study was located in the north of the province of Navarre (Northern Spain, see Fig. 1). In the territory the beechwood is well represented and corresponds to the south-western limit of the natural distribution area of European beech. According to the biogeographical areas and substrate, the principal beechwood types were selected:




Fig. 1. Map of the Iberian Peninsula and the location of 90 sampled plots in Navarre. The numbers correspond to the three principal beechwood types: 1 – *Carici sylvaticae* – *Fago sylvaticae* S.; 2 – *Saxifrago hirsutae* – *Fago sylvaticae* S.; 3 – *Scillo* – *Fago sylvaticae* S.

1) basophilous and ombrophilous beechwoods in the Cantabrian-Atlantic Province with calcareous soil (*Carici sylvaticae* – *Fago sylvaticae* S.); 2) acidophilous and ombrophilous ones in the Cantabrian-Atlantic Province with acidic soil (*Saxifraga hirsutae* – *Fago sylvaticae* S.); 3) Pyrenean basophilous and ombrophilous beechwoods in the Pyrenean Province with calcareous soil (*Scillo* – *Fago sylvaticae* S.) (Loidi, Báscos, 1995). Climatic conditions and vegetation vary considerably among the classified forest stands (Table 1). Pyrenean ones (beechwood type 3) have a more mountainous character than others, growing at higher altitude and with lower temperatures. In the Cantabrian-Atlantic region two beechwood types can be differentiated according to the substratum and rainfall characteristics: on the one hand, beechwoods with basophilous substratum and with low precipitation (beechwood type 1), and, on the other hand, those growing on acidophilous substratum and with higher precipitation (beechwood type 2). All stands are dominated by beech, but in some cases other tree species also occur (Table 1).

## DATA COLLECTION

According to three different beechwood types, a stratified random sampling scheme was used for data collection. In each beechwood type, 30 rectangular plots of 100 m<sup>2</sup> were randomly located

(Fig. 1). To ensure the presence of all woody debris types, the randomization was restricted by the presence of at least one big log. For each plot all woody debris were checked for fungal identification and all lignicolous macromycetes were identified (fungi with sporocarps larger than 1 mm), registering their frequency for different woody debris types according to the diameter, decay stage and host species. When microscopic identification was necessary the material was removed for further identification in the laboratory.

According to the diameter, the following classification of woody debris has been used: 1) Very Fine Woody Debris (VFWD): branches and twigs with  $\leq 5$  cm diameter; 2) Fine Woody Debris (FWD): logs with diameter between 5 and 10 cm; 3) Coarse Woody Debris (CWD): logs or snags with diameter  $\geq 10$  cm.

Decay stages were defined following a modified classification of Renvall (1995): (I) wood hard, trunk or branch is a solid piece, pushed knife penetrates only a few mm into the wood; (II) wood fairly hard, bark usually present but not firmly attached, pushed knife penetrates 1–2 cm into the wood; (III) wood fairly soft, small areas of sapwood already decomposed and without bark where knife penetrates easily; (IV) wood soft, wood pieces extensively decayed and usually large sections of the wood completely decomposed, the knife penetrates through the wood easily; (V) wood very soft, almost completely decomposed

Table 1. Principal characteristics of the sampled beechwoods

| Beechwood type               | Size (ha) | Other trees or shrubs   | Elevation range (m) | Substratum (Bedrock) | Tave <sup>1</sup> (°C) | Tmin <sup>2</sup> (°C) | Tmax <sup>3</sup> (°C) | Precipitation <sup>4</sup> (mm) | Ic <sup>5</sup> |
|------------------------------|-----------|---|---------------------|----------------------|------------------------|------------------------|------------------------|---------------------------------|-----------------|
| <b>Basophilous (Type 1)</b>  | 24.7      | <i>Crataegus monogyna</i> ,<br><i>C. laevigata</i>                            | 525–1 460           | limestone, dolomite  | 11                     | 4.5                    | 18.5                   | 1 300–1 600                     | 14              |
| <b>Acidophilous (Type 2)</b> | 24        | <i>Ilex aquifolium</i> , <i>Quercus robur</i>                                 | 225–1 550           | sandstone, clay      | 12                     | 5.5                    | 19                     | 2 000–2 500                     | 13              |
| <b>Pyrenean (Type 3)</b>     | 19        | <i>Abies alba</i> ,<br><i>Pinus sylvestris</i> ,<br><i>Buxus sempervirens</i> | 800–1 700           | limestone, sandstone | 9                      | 2                      | 16.5                   | 1 500                           | 15              |

<sup>1</sup> Mean annual temperature

<sup>2</sup> Mean temperature of the coldest month

<sup>3</sup> Mean temperature of the warmest month

<sup>4</sup> Annual precipitation

<sup>5</sup> Simple continentality index (mean annual temperature range)

Rivas-Martínez, Rivas-Sáenz (1996–2009)

and disintegrates easily between fingers. The decay stage was obtained for each fungal species, measuring in the area where fruitbodies grew.

The fieldwork was carried out in 2010 during the main period of fruiting production (end of September to early December) with one visit to each plot. A species found on a single piece of wood was considered as one record, regardless of the number of fruitbodies. The frequency of each species was recorded by the number of records in each woody debris piece of each plot.

The collected fruitbodies were identified, mostly following Eriksson, Ryvariden (1973, 1975, 1976); Eriksson et al. (1978, 1981, 1984); Hjortstam et al. (1987, 1988); Jülich, Stalpers (1980); Tellería, Melo (1995); Bernicchia (2005); Breitenbach, Kränzlin (1986); Bernicchia, Gorjón (2010). In addition, for some groups special literature was consulted (e. g. Boidin, 1994; Dueñas, 2005).

## DATA ANALYSIS

To analyze and compare fungal communities multivariate analyses were carried out using PRIMER-E and PERMANOVA+ for PRIMER 6 software package (Clarke, Gorley, 2006; Anderson et al., 2008). To observe the similarity of fungal communities Bray-Curtis resemblance index was used (Bray, Curtis, 1957). To determine the response of fungal composition to factors (e. g. woody debris class, beechwood type and wood decay stage) with the analysis of variance, Permanova statistical method was used. To observe the samples grouping in two-dimensional scale, canonical analysis of principal coordinates (CAP) was run. Spearman's rank coefficient was used to observe correlation between species and formed groups. Permutational tests of homogeneity of multivariate dispersions (PERMDISP) were used to check the homogeneity of groups.

## RESULTS

### Fungal diversity

In total, 3799 records of 274 species were collected over the period of investigation in all the study plots. The species belong to 143 genera, 25 orders and 2 phyla. The largest part of fungi belongs to the phylum *Basidiomycota* (90%), whereas the phylum *Ascomycota* comprises a smaller part (10%). The

number of species found in debris of each beechwood type were similar, 180 species in basophilous, 170 species in acidophilous and 169 in Pyrenean beechwoods. Five most common species of the phylum *Ascomycota* were *Diatrype disciformis* (Hoffm.) Fr. (344 records), *Hypoxylon fragiforme* (Pers.) J. Kickx f. (329 records), *Biscogniauxia nummularia* (Bull.) Kuntze (119 records), and *Kretzschmaria deusta* (Hoffm.) P. M. D. Martin (72 records); Five most common species of *Basidiomycota* were *Stereum ostrea* (Blume, T. Nees) Fr. (152 records), *Xenasmatella vaga* (Fr.) Stalpers (115 records), *Stereum hirsutum* (Willd.) Pers. (113 records), *Fomes fomentarius* (L.) J. J. Kickx (112 records) and *Trametes gibbosa* (Pers.) Fr. (95 records). On the other hand, 28% of the collected species are rare, with just a single record.

Fig. 2 shows the distribution of species richness and number overlap in three woody debris classes. As shown in this figure, the number of species changed with woody debris size, and the largest part of the species (201 species) appeared on CWD, while on FWD were found 164, and on VFWD 157. Moreover, CWD being the most diverse woody debris type in the number of species, it also had the majority of exclusive species (70). The number of species that different woody debris classes shared also varied and debris types that share the highest number of species were CWD and FWD, while debris with lower species in common were CWD and VFWD.





Fig. 2. Species richness and number overlap in three woody debris classes



Although most lignicolous species were found on CWD, not all the fungal groups had shown the same trend (Fig. 3). In case of Ascomycota, 50% of species were found exclusively on VFWD, 20% exclusively on FWD and 30% exclusively on CWD, whereas Basidiomycota had shown an inverse distribution among each woody debris class (Fig. 3).

The greatest part of fungi (69%) was found in the earliest decay stages (I and II), whereas in advanced decayed wood the number of species found was lower, 48% in III decay stage and 37% in IV and V decay stages.




**Fig. 3.** The proportion of the number of species found exclusively on each woody debris class (VFWD, FWD and CWD) for Basidiomycota and Ascomycota

## Fungal community

The differences in species composition among different woody debris classes of sampled plots are precisely visualized in the CAP analysis (Fig. 4). The canonical analysis of principal coordinates revealed that fungal communities from different woody debris classes were statistically different (Permanova test,  $p = 0.0001$ ) and that fungal composition changed gradually, from smaller diameter woody debris towards coarser ones, and vice versa (axis  $x$ ). The correlation between axis  $x$  and fungal composition was 0.69. This analysis also demonstrated that fungal communities were different in faithful species, VFWD and CWD having a higher number of constant and frequent species (axis  $y$ ), like *Diatrype disciformis* (Hoffm.) Fr. and *Bisporella citrina* (Batsch) Korf, S. E. Carp. on VFWD and *Fomes fomentarius* (L.) J. J. Kickx and *Kretzschmaria deusta* (Hoffm.) P. M. D. Martin on CWD (species with at least 0.5 Spearman's correlation value), whilst FWD was characterized by the absence of such faithful species (Fig. 4, Table 2).

Permanova test showed that there was an interaction between beechwood type and woody debris class ( $p = 0.0051$ ). In this way fungal communities



**Fig. 4.** Canonical analysis of principal coordinates (CAP) for three woody debris classes of the sampled plots. Species with at least 0.5 Spearman's correlation value are also given

Table 2. List of faithful species (species having Spearman's correlation value  $\geq 0.25$ ) among groups from the CAP analysis (VFWD, FWD and CWD)

| CWD                          | Spearman's correlation | VFWD                             | Spearman's correlation |
|------------------------------|------------------------|----------------------------------|------------------------|
| <i>Fomes fomentarius</i>     | 0.75                   | <i>Diatrype disciformis</i>      | 0.8                    |
| <i>Kretzschmaria deusta</i>  | 0.6                    | <i>Bisporella citrina</i>        | 0.5                    |
| <i>Fomitopsis pinicola</i>   | 0.4                    | <i>Gloeocystidiellum porosum</i> | 0.4                    |
| <i>Oudemansiella mucida</i>  | 0.35                   | <i>Exidia thuretiana</i>         | 0.4                    |
| <i>Ascocoryne cylichnium</i> | 0.25                   | <i>Peniophora cinerea</i>        | 0.3                    |
| <i>Ganoderma pfeifferi</i>   | 0.25                   | <i>Athelopsis glaucina</i>       | 0.25                   |
| <i>Trichaptum biforme</i>    | 0.25                   | <i>Skeletocutis nivea</i>        | 0.25                   |

of each woody debris class in each beechwood type were different. In the analysis of fungal composition for each forest type it has been observed that these were dissimilar, with one exception, fungal communities of CWD, in Pyrenean and basophilous beechwoods (Pair Wise test). In Fig. 5 the spa-

tial ordination of beechwood plots for each woody debris class is represented. The most associated species of acidophilous beechwoods were *Trametes versicolor* (L.) Lloyd on all woody debris classes, *Hyphoderma setigerum* (Fr.) Donk on FWD, and *Trichaptum biforme* (Fr.) Ryvarden on FWD and




Fig. 5. Canonical analysis of principal coordinates (CAP) in three woody debris classes for all beechwood plots. Species with at least 0.4 Spearman's correlation value are also given


CWD. For basophilous ones the most associated species were *Exidia thurettiana* (Lév.) Fr. and *Diatrype disciformis* (Hoffm.) Fr. on VFWD. In case of Pyrenean beechwoods, the most associated species was *Royoporus badius* (Pers.) A. B. De on CWD. The species *Xenasmatella vaga* (Fr.) Stalpers was faithful in beechwoods with a more continental character (Pyrenean and basophilous ones) both on VFWD and FWD.

The structure and composition of fungal community was different according to wood decay stages (Fig. 6), and the CAP analysis revealed that there were three statistically different groups (Permanova test,  $p = 0.0001$ ). Firstly, the fungal communities of the earliest stages of decomposition process (I and II decay stages), followed by communities of intermediate decomposition stages (III decay stage), and finally the communities that grow in strongly decayed wood (IV and V decay stages).

## DISCUSSION

In this study, fungal diversity has been quantified recording the presence of macromycetes on wood. Although there were certainly mycelia which did not produce fruitbodies, the macromycetes inventory is presently assumed as one of the most appropriate methods in large field surveys, because molecular methods are still significantly skewed (Avis et al., 2010), expensive and not yet available for many species (Ovaskainen et al., 2010); whilst fruitbody inventory provides data suitable for comparison between different woody debris classes (Heilmann-Clausen, Christensen, 2004; Küffer, Senn-Irlet, 2005b).

The properties of dead wood vary depending on woody debris size and decomposing process. On the one hand, woody debris diameter determines the microclimate, decomposition rate, ecological niches diversity and the amount of



**Fig. 6.** Canonical analysis of principal coordinates (CAP) for the principal degrees of decomposition of the sampled plots

substrate. On the other hand, in the course of decomposition process, woody debris undergoes structural and chemical changes. The effect of these two variables on fungal communities composition is shown in this study by the CAP analyses.

In terms of woody debris diameter, our results have shown that fungal composition changes significantly in three woody debris classes, and that each debris category possesses species which grow exclusively on it. Thereby, fungal communities that grow in VFWD must be specialized to unstable climatic conditions, high decomposition rates, and high surface area and wood volume ratio (i. e. less wood volume per surface area unit). Nevertheless, CWD offers to wood inhabiting fungi a more stable resource and a large amount of substrate. In this way, both VFWD and CWD are characterized by constant and frequent species such as *Bisporella citrina* and *Diatrype disciformis* on VFWD, and *Fomes fomentarius* and *Kretzschmaria deusta* on CWD (with a high correlation value between debris class and species). The conditions that FWD offers are between those given by VFWD and CWD remains. Thus the fungal community found in this woody debris class, in many cases, overlapped the other two communities and is marked by the absence of faithful species.

Woody debris with diameter less than 10 cm has often been neglected in many earlier studies, and only CWD has been studied, due to CWD volume in managed forests being considerably lower than in unmanaged ones, and also because wood inhabiting organisms diversity per unit is higher on CWD than on thinner ones. However, the interest in understanding mycobiota diversity related to thinner woody debris has recently increased, after VFWD class had been proposed. Although the number of exclusive species decreases in thinner woody debris, it is remarkable that VFWD is especially important for specific groups of fungi, such as Ascomycota, which are adapted to live in branches and twigs (Rayner, Boddy, 1997). Nordén et al. (2004) were the first to point out the difference in frequency between Ascomycota and Basidiomycota on CWD and FWD. The data of this study are in agreement with the results obtained by Nordén et al. in deciduous forests of Central

Europe, where the importance of small diameter woody debris for the phylum *Ascomycota* was revealed.

Several studies have shown that fungal community changes during wood decomposition process (Renvall, 1995; Heilmann-Clausen, Christensen, 2003; Heilmann-Clausen et al., 2005), and some authors hold that decay gradient is the dominant factor structuring communities (Høiland, Bendiksen, 1996; Lindblad, 1997). As can be seen in Fig. 6, according to wood decay stage three main groups can be distinguished: (1) pioneer species, preferring recently fallen or weakly decomposed wood, (2) intermediate stage species, which replace previous species when easily decomposable components of the wood were consumed, (3) late stage species, the ones that are specialised in degrading more complex components of the wood. This succession is often very strict, and in some cases the presence of one species depends on the preceding one (Renvall, 1995; Heilmann-Clausen, Boddy, 2005).

Contrary to the results in other studies of Europe (e. g. Renvall, 1995), in our study the majority of species were found on wood of early decomposition stages, and the lowest number of species was found on strongly decayed wood. However, this can be explained by the fact that the largest part of the sampled beechwood area is currently or recently managed, so most of the examined woody debris was in the earliest decay stage and fungal communities from the latest decay stages are not completely developed.

An overall change of fungal composition has been observed along the climatic gradient from acidophilous beechwoods with an oceanic character, to Pyrenean and basophilous beechwoods with a more continental character. This general pattern is repeated in the analysis of species composition among woody debris types, and consequently faithful species of each woody debris type have varied according to the climatic characteristics.

## ACKNOWLEDGEMENTS

We are grateful to G. García-Baquero for his valuable comments on methodological design. We thank the government of Navarre for making fieldwork easier providing permission when it was needed. We are also indebted to the University of



the Basque Country (UPV/EHU) for the given fellowship (PIF10/2010/PIF10008) that has partially supported this work. We also want to thank two anonymous reviewers for critically reviewing the manuscript.

Received 29 September 2011

Accepted 21 November 2011

## REFERENCES

1. Anderson M. J., Gorley R. N., Clarke K. R. 2008. PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods. Plymouth, UK.
2. Avis P. G., Branco S., Tang Y., Mueller G. M. 2010. Pooled samples bias fungal communities descriptions. *Molecular Ecology Resources*. Vol. 10: 135–141.
3. Bader P., Jansson S., Jonsson B. G. 1995. Wood-inhabiting fungi and substratum decline in selectively logged boreal spruce forests. *Biological Conservation*. Vol. 72: 355–362.
4. Bernicchia A. 2005. *Fungi Europaei: Polyporaceae s.l.* Alassio: Edizioni Candusso.
5. Bernicchia A., Gorjón S. P. 2010. *Fungi Europaei: Corticiaceae s.l.* Alassio: Edizioni Candusso.
6. Boddy L. Development and Function of Fungal Communities in Decomposing Wood. In: Carrol G. C. and Wicklow D. T. (eds.), *The Fungal Community, Its Organization and Role in the Ecosystem*, 2nd ed. New York: Marcel Dekker. 1992: 749–782.
7. Boddy L. 2001. Fungal community ecology and wood decomposition processes in angiosperms: from standing tree to complete decay of coarse woody debris. *Ecological Bulletins*. Vol. 49: 43–53.
8. Boidin J. 1994. Les Peniophoraceae des parties tempérées et froides de l'hémisphère nord (Basidiomycotina). *Bulletin mensuel de la Société Linnée de Lyon*. Vol. 63: 317–334.
9. Bray J. R., Curtis J. T. 1957. An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs*. Vol. 27: 325–349.
10. Breitenbach J., Kränzlin F. *Fungi of Switzerland II (Heterobasidiomycetes, Aphyllophorales, Gastromycetes)*. Luzern: Verlag Mykologia; 1986.
11. Clarke K. R., Gorley R. N. 2006. PRIMER v6: User manual / tutorial, PRIMER-E, Plymouth.
12. Dueñas M. 2005. New and interesting Iberian heterobasidiomycetous fungi I. *Nova Hedwigia*. Vol. 81(1–2): 177–198.
13. ECCB (European Committee for Conservation of Bryophytes). *Red Data Book of European Bryophytes*. Cramer, J. Berlin; 1995.
14. Eriksson J., Hjortstam K., Ryvarden L. 1978. The Corticiaceae of North Europe (Mycoaciella-Phanerochaete). *Fungiflora (Oslo)*. Vol. 5: 887–1048.
15. Eriksson J., Hjortstam K., Ryvarden L. 1981. The Corticiaceae of North Europe (Phlebia-Sarcodontia). *Fungiflora (Oslo)*. Vol. 6: 1049–1276.
16. Eriksson J., Hjortstam K., Ryvarden L. 1984. The Corticiaceae of North Europe (Schizopora-Suillosporium). *Fungiflora (Oslo)*. Vol. 7: 1279–1449.
17. Eriksson J., Ryvarden L. 1973. The Corticiaceae of North Europe Vol (Aleurodiscus-Confertobasidium). *Fungiflora (Oslo)*. Vol. 2: 59–286.
18. Eriksson J., Ryvarden L. 1975. The Corticiaceae of North Europe Vol (Coronicium-Hyphoderma). *Fungiflora (Oslo)*. Vol. 3: 287–546.
19. Eriksson J., Ryvarden L. 1976. The Corticiaceae of North Europe (Hyphodermella-Mycoacia). *Fungiflora (Oslo)*. Vol. 4: 574–886.
20. Harmon M. E., Franklin J. F., Swanson F. J., Sollins P., Gregory S. V., Lattin J. D., Anderson N. H., Cline S. P., Aumen N. G., Sedell J. R., Lienkaemper G. W., Cromack K., Cummins K. W. 1986. Ecology of Coarse Woody Debris in Temperate Ecosystems. *Advances in Ecological Research*. Vol. 15: 133–276.
21. Heilmann-Clausen J. 2001. A gradient analysis of communities of macrofungi and slime moulds on decaying beech logs. *Mycological Research*. Vol. 105: 575–596.
22. Heilmann-Clausen J., Aude E., Christensen M. 2005. Cryptogam communities on decaying deciduous wood – does tree species diversity matter? *Biodiversity and Conservation*. Vol. 14: 2061–2078.
23. Heilmann-Clausen J., Boddy L. 2005. Inhibition and stimulation effects in communities of wood decay fungi: exudates from colonized wood influence growth by other species. *Microbial Ecology*. Vol. 49: 399–406.
24. Heilmann-Clausen J., Christensen M. 2003. Fungal diversity on decaying beech logs – implications for sustainable forestry. *Biodiversity and Conservation*. Vol. 12: 953–973.
25. Heilmann-Clausen J., Christensen M. 2004. Does size matter? On the importance of various dead

- wood fractions for fungal diversity in Danish beech forests. *Forest Ecology and Management*. Vol. 201: 105–117.
26. Heilmann-Clausen J., Walley R. 2007. Some records of wood-inhabiting fungi on *Fagus sylvatica* in Northern Spain. *Revista Catalana de Micologia*. Vol. 29: 67–80.
  27. Hjortstam K., Larsson K., Ryvarden L. 1988. The Corticiaceae of North Europe (*Thanatephorus-Ypsilonidum*). *Fungiflora* (Oslo). Vol. 8: 1450–1631.
  28. Hjortstam K., Larsson K. H., Ryvarden L. 1987. The Corticiaceae of North Europe (Introduction and keys). *Fungiflora* (Oslo). Vol. 1: 1–59.
  29. Høiland K., Bendiksen E. 1996. Biodiversity of wood-inhabiting fungi in a boreal coniferous forest in Ser-Trendelag County, Central Norway. *Nordic Journal of Botany*. Vol. 16: 643–659.
  30. Jülich W., Stalpers J. A. The resupinate non-poroid Aphylllophorales of the temperate northern hemisphere. Amsterdam, Oxford, New York: North-Holland Publishing Company; 1980.
  31. Küffer N. 2008. Wood-inhabiting aphylllophoroid basidiomycetes: Diversity, Ecology and Conservation. Ph. D. Thesis, Neuchâtel: Université de Neuchâtel.
  32. Küffer N., Senn-Irlet B. 2005a. Influence of forest management on the species richness and composition of wood-inhabiting basidiomycetes in Swiss forests. *Biodiversity and Conservation*. Vol. 14: 2419–2435.
  33. Küffer N., Senn-Irlet B. 2005b. Diversity and ecology of wood inhabiting aphylllophoroid basidiomycetes on fallen woody debris in various forest types in Switzerland. *Mycological Progress*. Vol. 4: 77–86.
  34. Lindblad I. 1998. Wood-inhabiting fungi on fallen logs of Norway spruce: relations to forest management and substrate quality. *Nordic Journal of Botany*. Vol. 18: 243–255.
  35. Loidi J., Bascónes J. C. 1995. Memoria del mapa de series de vegetación de Navarra, E. 1:200.000. Pamplona: Dpto. de Medio Ambiente, Ordenación del Territorio y Vivienda, Gobierno de Navarra.
  36. Lonsdale D., Pautasso M., Holdenrieder O. 2008. Wood-decaying fungi in the forest: conservation needs and management options. *European Journal of Forest Research*. Vol. 127: 1–22.
  37. Niemelä T., Renvall P., Penttillä R. 1995. Interactions of fungi at late stages of wood decomposition. *Annales Botanici Fennici*. Vol. 32: 141–152.
  38. Nordén B., Ryberg M., Götmark F., Olausson B. 2004. Relative importance of coarse and fine woody debris for the diversity of wood inhabiting fungi in temperate broadleaf forests. *Biological Conservation*. Vol. 101: 1–8.
  39. Ovaskainen O., Nokso-Koivisto J., Hottola J., Rajala T., Pennanen T., Ali-Kovero H., Miettinen O., Oinonen P., Auvinen P., Paulin L., Larsson K.-H., Mäkipää R. 2010. Identifying wood-inhabiting fungi with 454 sequencing – What is the probability that BLAST gives the correct species? *Fungal Ecology*. Vol. 3: 274–283.
  40. Peterken G. F. *Natural Woodland: Ecology and Conservation in Northern Temperate Regions*. Cambridge: Cambridge University Press; 1996.
  41. Rayner A. D. M., Boddy L. *Fungal Decomposition of Wood: Its Biology and Ecology*. Bath: Bath Press; 1997.
  42. Renvall P. 1995. **Community structure and dynamics** of wood-rooting Basidiomycetes on decomposing conifer trunks in Northern Finland. *Karstenia*. Vol. 3: 1–51.
  43. Rivas-Martínez S., Rivas-Sáenz S. 1996–2009. *Worldwide Bioclimatic Classification System*. Phytosociological Research Center, Spain. <http://www.globalbioclimatics.org>.
  43. Siitonen J. 2001. Forest management, coarse woody debris and saproxylic organisms: fennoscandian boreal forests as an example. *Ecological Bulletins*. Vol. 49: 11–41.
  45. Sippola A. L., Renvall P. 1999. Wood-decomposing fungi and seed-tree cutting: A 40 year perspective. *Forest Ecology and Management*. Vol. 115: 183–201.
  46. Swift M. J. Basidiomycetes as components of forest ecosystems. In: Frankland J. N., Hedger J. N. and Swift M. J. (eds.), **Decomposer basidiomycetes – their biology and ecology** Cambridge: Cambridge University Press; 1982: 307–337.
  47. Tellería M. T. Riqueza fúngica de la Península Ibérica e Islas Baleares: El proyecto “Flora Micológica Ibérica”. In: Pineda F. D., Casado M. A., de Miguel J. M., Montalvo J. (eds.), *La diversidad biológica de España*. Madrid: Prentice Hall; 2002: 153–160.
  48. Tellería M. T., Melo I. Aphylllophorales resupinate non poroides, I (*Acanthobasidium-Cystostereum*). Madrid: CSIC – Real Jardín Botánico; 1995.

Nerea Abrego, Isabel Salcedo

## GRYBŲ ĮVAIROVĖS KAITA REMIANTIS MEDIENOS LIEKANŲ TIPU (ŠIAURĖS ISPANIJOS TYRIMŲ PAVYZDYS)

### *Santrauka*

Lignofilinių grybų bendrijų tyrimai Pietų Europoje yra nepakankami ir dauguma jų atlikti tiriant stambias medienos liekanas (SML). Plonos (PML) ir labai plonos medienos liekanos (LPML) tiriamos retai. Ieškant lignofilinių makroskopinių grybų buvo patikrintos visos medienos liekanos iš 90-ies buko miškų plotelių (Navara, Šiaurės Ispanija). Nors daugelis rūšių buvo rasta ant SML, mažo dydžio medienos liekanos pasižymėjo *Ascomycota* rūšių įvairove, o didelio dydžio medienos liekanose rasta *Basidiomycota* rūšių. Kanoninė analizė (CAP) atskleidė, kad grybų bendrijos ant skirtingo skersmens medienos liekanų yra nepanašios ir kad grybų sudėtis palaipsniui keičiasi pereinant nuo mažesnio skersmens medienos liekanų prie stambesniųjų ar atvirkščiai. Susiformavusios grupės taip pat skiriasi grybų rūšių įvairove – LPML ir SML grupėse buvo daugiau dažnesnių rūšių. Nustatyta grybų sudėties kaitos priklausomybė nuo buko miško tipo; taigi buvo rastos kiekvienam medienos liekanų tipui skirtinguose buko miškuose būdingos rūšys. Grybų bendrijos struktūra ir sudėtis kito skirtingose irimo stadijose, tad kanoninė analizė leido išskirti tris pagrindines grupes: bendrijas, susijusias su silpnai, vidutiniškai ir stipriai suirusia mediena.

**Raktažodžiai:** irimo stadija, diametras, *Fagus sylvatica*, lignofilinių grybų bendrija, makroskopiniai grybai, Navara

